



Contents lists available at ScienceDirect

## Global and Planetary Change

journal homepage: [www.elsevier.com/locate/gloplacha](http://www.elsevier.com/locate/gloplacha)

## Comment on “Scrutinizing the carbon cycle and CO<sub>2</sub> residence time in the atmosphere” by H. Harde

Peter Köhler<sup>a,\*</sup>, Judith Hauck<sup>a</sup>, Christoph Völker<sup>a</sup>, Dieter A. Wolf-Gladrow<sup>a</sup>, Martin Butzin<sup>a</sup>, Joshua B. Halpern<sup>b</sup>, Ken Rice<sup>c</sup>, Richard E. Zeebe<sup>d</sup>

<sup>a</sup> Alfred-Wegener-Institut Helmholtz-Zentrum für Polar-und Meeresforschung (AWI), P.O. Box 12 01 61, Bremerhaven 27515, Germany

<sup>b</sup> Department of Chemistry, Howard University, Washington, DC 20059, USA

<sup>c</sup> SUPA, Institute for Astronomy, University of Edinburgh, The Royal Observatory, Blackford Hill, Edinburgh EH93HJ, UK

<sup>d</sup> SOEST, University of Hawaii at Manoa, Honolulu, HI 96822, USA

## ARTICLE INFO

## Keywords:

Carbon cycle

Anthropogenic carbon

## ABSTRACT

Harde (2017) proposes an *alternative accounting scheme* for the modern carbon cycle and concludes that only 4.3% of today's atmospheric CO<sub>2</sub> is a result of anthropogenic emissions. As we will show, this *alternative scheme* is too simple, is based on invalid assumptions, and does not address many of the key processes involved in the global carbon cycle that are important on the timescale of interest. Harde (2017) therefore reaches an incorrect conclusion about the role of anthropogenic CO<sub>2</sub> emissions. Harde (2017) tries to explain changes in atmospheric CO<sub>2</sub> concentration with a single equation, while the most simple model of the carbon cycle must at minimum contain equations of at least two reservoirs (the atmosphere and the surface ocean), which are solved simultaneously. A single equation is fundamentally at odds with basic theory and observations. In the following we will (i) clarify the difference between CO<sub>2</sub> atmospheric residence time and adjustment time, (ii) present recently published information about anthropogenic carbon, (iii) present details about the processes that are missing in Harde (2017), (iv) briefly discuss shortcoming in Harde's generalization to paleo timescales, (v) and comment on deficiencies in some of the literature cited in Harde (2017).

### 1. Residence time versus adjustment time

The global carbon cycle is currently not in a steady state as shown, for example, by measurements of atmospheric CO<sub>2</sub> concentration at Mauna Loa (Hawaii) and at the South Pole since 1958 (Dlugokencky et al., 2016). The main reason for this increase is the addition of ‘anthropogenic CO<sub>2</sub>’ by burning of coal, oil, and gas, industrial processes and land use change (Le Quéré et al., 2016). In the case of non-steady state conditions one can ask the question: How long will the perturbation (here: higher atmospheric CO<sub>2</sub> concentration) stay? If one adds a certain amount of anthropogenic CO<sub>2</sub> to the atmosphere at time  $t_0$ , the concentration will increase suddenly and then fall off following a complicated function that depends on the response of the various active carbon reservoirs (surface ocean, intermediate and deep ocean, marine sediments, terrestrial biosphere). The time connected to such a relaxation in atmospheric CO<sub>2</sub> concentration is the **adjustment (or equilibration) time** and the timescale of interest for the problem at hand. The function how CO<sub>2</sub> relaxes after such an initial perturbation can be approximated by the sum of a few exponential functions with

different characteristic timescales (e.g. Archer et al., 1997; Lord et al., 2016). Simple one-box models suggest adjustment times of about 70 years, but these models ignore many relevant processes and consequently underestimate this timescale (Cawley, 2011). More complex models suggest adjustment (equilibration) times of well over 100 years, and that it depends on total anthropogenic emissions (Archer et al., 2009; Joos et al., 2013; Lord et al., 2016). When anthropogenic CO<sub>2</sub> is added continuously one has to run a global carbon cycle model that takes into account the responses by the various reservoirs mentioned above (e.g. Friedlingstein et al., 2006; Jones et al., 2013).

The timescale determined in Eqs.(7) and (8) in Harde (2017) is actually an approximation of the **residence time**, i.e. the average length of time for which an individual molecule of CO<sub>2</sub> remains in the atmosphere before being taken up by the ocean or terrestrial biosphere. Given the fluxes into, and out of, the atmosphere, we would expect a CO<sub>2</sub> molecule to only remain in the atmosphere for a few years, before being replaced by a molecule from one of the other reservoirs. The usual misunderstanding is that this is CO<sub>2</sub> leaving the atmosphere, rather than mostly just being exchanged, leaving no change in

\* Corresponding author.

E-mail address: [Peter.Koehler@awi.de](mailto:Peter.Koehler@awi.de) (P. Köhler).

<http://dx.doi.org/10.1016/j.gloplacha.2017.09.015>

Received 24 March 2017; Received in revised form 22 September 2017; Accepted 22 September 2017

0921-8181/ © 2017 Elsevier B.V. All rights reserved.

atmospheric CO<sub>2</sub> concentration. Even though the numbers presented in Harde (2017) are reasonable approximations for the residence time, they are largely irrelevant for what the paper later presents.

Changes in atmospheric CO<sub>2</sub> concentration following the initial perturbation depend on the net CO<sub>2</sub> flux out of the atmosphere, rather than - as in the case of the residence time - depending only on the flux into the natural sinks. Note that the residence and adjustment times refer to different and distinct aspects of the carbon cycle and have different definitions; a distinction clearly made in the IPCC First Assessment Report (Houghton et al., 1990, §1.2.1) as well as in more recent reports (Stocker et al., 2013, p. 1457). Thus to conflate residence time and adjustment time is a fundamental misunderstanding of the carbon cycle.

Given this difference between the residence time (years) and adjustment time (centuries to millennia) we would also not expect an enhancement in atmospheric CO<sub>2</sub> to be entirely composed of molecules of directly anthropogenic origin, even if the cause for such an enhancement were entirely anthropogenic. Therefore, the claim in Harde (2017) that the anthropogenic contribution makes up only 15% of the increase since the industrial era - even if correct - is not an indication that the increase is not entirely anthropogenic.

## 2. Most recent anthropogenic carbon inventory

Total anthropogenic CO<sub>2</sub> from fossil fuel and cement production emitted between 1750 and 2010 has accumulated to 365 PgC (Le Quéré et al., 2016). A further 153 PgC was emitted in the same time interval from land use changes (Le Quéré et al., 2016). In 2010 the atmospheric CO<sub>2</sub> concentration was approximately 390 ppm (Dlugokencky et al., 2016), a value that features prominently in the calculations of Harde (2017).

Anthropogenic carbon in the ocean can be tracked by various methods, e.g. the <sup>13</sup>C Suess effect (Eide et al., 2017), ΔC\* method (Gruber et al., 1996), or anthropogenic produced substances, such as CFCs (Lauvset et al., 2016). Various approaches have shown that the oceanic sink accounts for 48% of the total fossil-fuel and cement-manufacturing CO<sub>2</sub> emissions of 118 PgC emitted between 1800 and 1994 (Sabine et al., 2004). Landschützer et al. (2016) calculated an increase in anthropogenic carbon in the ocean of about 60 PgC released by fossil-fuel and cement-manufacturing CO<sub>2</sub> emissions between 1982 and 2012. Taking the temporal overlay of both studies into account, we find an anthropogenic carbon inventory of the fossil-fuel and cement-manufacturing CO<sub>2</sub> emissions 1800–2012 in the ocean of about 150 PgC. This is 41% of the accumulated emission from fossil fuel and cement production, or 29% of the total emissions including land use change. In this scenario, the rise in atmospheric CO<sub>2</sub> from a pre-industrial value of 278 ppm (= 589 PgC) before 1750, to 390 ppm (= 827 PgC) in 2010 is solely due to anthropogenic emissions. They overprint any potential natural CO<sub>2</sub> outgassing from the ocean (see the decomposition of anthropogenic and natural fluxes between ocean and atmosphere in Gruber et al. (2009)). The rise in the atmospheric carbon reservoir by 112 ppm, or 238 PgC, corresponds to an airborne fraction of 46% of the total anthropogenic emissions of 518 PgC. The missing residual of the anthropogenic emissions of 130 PgC is assigned to terrestrial carbon uptake.

According to Harde (2017), 4.3% of the actual atmospheric CO<sub>2</sub> concentration is of anthropogenic origin. With an atmospheric CO<sub>2</sub> concentration of 390 ppm used in Harde (2017), reached in ~2010, this is similar to a proposed anthropogenic CO<sub>2</sub> concentration of about 17 ppm or 36 PgC. It would imply that only 7% of the total anthropogenic emissions remained airborne. The airborne fraction of Harde (2017) is therefore a factor of 6.6 smaller than in the inventory that is supported by observational-based studies. If the approach in Harde (2017) was correct, it would directly asked for evidence where this anthropogenic carbon has been stored. Unfortunately, no further evidence for this storage has been given in the paper and as we have shown

above, it cannot reside in the ocean.

## 3. Why is the Harde model too simple?

The core argument in Harde (2017), section 3, is about the lifetime of anthropogenic CO<sub>2</sub> in the atmosphere, closely related to the airborne fraction that remains after a given time. A framework is then developed in which both natural and anthropogenic carbon fluxes are analyzed. In this framework one important part of the carbon cycle, which is of major relevance for the airborne fraction of CO<sub>2</sub>, is missing: the carbonate chemistry in the ocean. It is correctly stated that Henry's Law governs the net gas exchange of CO<sub>2</sub> between the surface ocean and the atmosphere, with higher temperatures leading to a higher net flux to the atmosphere. However, within the ocean CO<sub>2</sub> molecules react with water to form carbonic acid (H<sub>2</sub>CO<sub>3</sub>) which subsequently dissociates into hydrogen ions (H<sup>+</sup>) and bicarbonate ions (HCO<sub>3</sub><sup>-</sup>). In a second step the HCO<sub>3</sub><sup>-</sup>-ion dissociates into another H<sup>+</sup>-ion and a carbonate ion (CO<sub>3</sub><sup>2-</sup>). The sum of all these carbon species is typically referred to as dissolved inorganic carbon (DIC). For present day conditions the fraction of carbonic acid is negligible; the majority of DIC (~90%) is found as HCO<sub>3</sub><sup>-</sup>, ~9% as CO<sub>3</sub><sup>2-</sup>, and only about 1% is found as dissolved CO<sub>2</sub> (Zeebe and Wolf-Gladrow, 2001). Only this 1% of DIC in the surface ocean, found as dissolved CO<sub>2</sub>, can exchange with the atmosphere. Thus, the carbonate chemistry represents a bottleneck for the oceanic uptake of anthropogenic CO<sub>2</sub> emitted to the atmosphere. Note, that the basic knowledge on the marine carbonate system, which is completely neglected in Harde (2017), is at least 60 years old, e.g. see Revelle and Suess (1957) and references therein. Furthermore, different software packages to compute the marine carbonate chemistry have been published in the meantime (e.g. Orr et al., 2015), and are in most cases freely available, e.g. see <http://cdiac.ornl.gov/ftp/oceans/co2sys/> for different versions of the package CO2SYS which was widely discussed in (e.g. Orr et al., 2015) or [https://www.soest.hawaii.edu/oceanography/faculty/zeebe\\_files/CO2\\_System\\_in\\_Seawater/csys.html](https://www.soest.hawaii.edu/oceanography/faculty/zeebe_files/CO2_System_in_Seawater/csys.html) for Matlab routines to Zeebe and Wolf-Gladrow (2001).

This effect of the carbonate chemistry on the carbon cycle is not a theoretical concept, but an observed quantity also known as the Revelle (or buffer) factor *R*. This is a fundamental property of the marine carbonate system and is implicitly considered in marine carbon cycle models underlying the analyses summarized in the IPCC-AR5, the 5th Assessment Report of the Intergovernmental Panel on Climate Change (Stocker et al., 2013). The carbonate chemistry in seawater describing these processes in detail is well known (compare, for example Dickson et al., 2007; Zeebe and Wolf-Gladrow, 2001). The Revelle factor is defined as the ratio of the relative change of dissolved CO<sub>2</sub> to the relative change of DIC and can be readily calculated:

$$R = \frac{\Delta\text{CO}_2/\text{CO}_2}{\Delta\text{DIC}/\text{DIC}} \quad (1)$$

From open ocean data it is known that *R* varies between 8 and 15 (Sabine et al., 2004). A Revelle factor of 8, for example, leads to a DIC increase by only 12.5% for a doubling of dissolved CO<sub>2</sub>. A rise in atmospheric and oceanic carbon content goes along with an increase in the Revelle factor, a phenomenon which is already measurable (e.g. Hauck et al., 2010). This implies that the oceanic uptake of anthropogenic carbon will become slower if we continue to increase anthropogenic CO<sub>2</sub> emissions. This is already seen in all CMIP5 model simulations (Jones et al., 2013). The scientific literature describing the marine carbonate chemistry, which, if complete, automatically includes the Revelle effect, is based on decades of laboratory experiments, field observations and theoretical understanding of the underlying chemical processes and is very well established. The books by Dickson et al. (2007), Zeebe and Wolf-Gladrow (2001) are only two examples of the state of knowledge in this field.

The carbonate chemistry is the most relevant part of the carbon cycle, which is of importance on the timescale of interest, yet ignored in

the erroneous approach of Harde (2017). The uptake of anthropogenic CO<sub>2</sub> by the terrestrial part of the carbon cycle is also relevant (e.g. Joos et al., 2013), and nowadays approximately of similar size as the marine carbon uptake (Le Quéré et al., 2016). However, terrestrial carbon uptakes are in the global carbon budgets of Le Quéré et al. (2016) still estimated from the unexplained residual, after fossil fuels and emissions from industry and land-use change have been balanced by constraints on changes in carbon pools in the atmosphere and the ocean. We therefore refrain from a more in-depth discussion of terrestrial carbon uptake processes here. Further processes with different impact on the airborne fraction of anthropogenic CO<sub>2</sub>, which are of relevance if longer timescales are of interest (e.g. necessary for the generalization and application to the paleo data) are ocean overturning, carbonate compensation and continental weathering rates (Lord et al., 2016). If implemented in a model this results in an airborne fraction of anthropogenic CO<sub>2</sub> emissions of around 40% on a 100-year timescale falling to ~18% on a 1000-year timescale (Joos et al., 2013) and down to 5% and 2% on timescales of 10<sup>5</sup> and 10<sup>6</sup> years, respectively (Lord et al., 2016).

Remark: Most of these details above on the role of the carbonate chemistry have been taken from another comment some of the authors published as part of the online discussion on another, overly simplistic, and therefore biased approach to explain the modern carbon cycle (<http://www.earth-syst-dynam-discuss.net/6/C813/2015/esdd-6-C813-2015.pdf>).

**Harde's flawed 1-box carbon cycle:** One key element of Harde's carbon cycle is the assumption of a simple absorption/decay process, which is unsuitable for the problem at hand. Harde's Eq. (11) reads:

$$\frac{dC_{\text{CO}_2}}{dt} = e_T - \alpha \cdot C_{\text{CO}_2} \quad (2)$$

where  $C_{\text{CO}_2}$  is the atmospheric CO<sub>2</sub> concentration,  $e_T$  is a total emission rate,  $\alpha = 1/\tau$  is an absorption efficiency, and  $\tau$  is Harde's CO<sub>2</sub> "lifetime". Thus, Harde assumes that  $C_{\text{CO}_2}$  can be predicted by solving only a single rate equation of the carbon cycle (other reservoirs may exist but their time evolution is ignored). However, at any given time  $t$ , the CO<sub>2</sub> fluxes into and out of the atmosphere depend on, for instance, the atmosphere-ocean disequilibrium, which in turn depends on simultaneous changes in ocean carbon inventory and seawater chemistry, as explained above. Thus, even the simplest carbon cycle model must at minimum comprise two boxes for atmosphere and ocean (including Revelle factor), whose equations are solved simultaneously. For investigations of timescales longer than centuries (e.g. in paleo applications as done in the generalization) processes which export carbon from the surface to the deep ocean (so-called carbon pumps, see Volk and Hoffert, 1985) also need to be taken into consideration, asking for at least another deep ocean box. Yet, Harde (2017) ignores this fact (and many others) that have been established in over 60 years of carbon-cycle research (a few examples include Revelle and Suess, 1957; Oeschger et al., 1975; Heimann and Maier-Reimer, 1996; Archer et al., 2009; Joos et al., 2013). As a result, the approach in Harde (2017) leads to fundamentally flawed mass balances, CO<sub>2</sub> "lifetimes", and thus erroneous conclusions.

Note also that the posited analogy to radiocarbon (and other isotopes) is incorrect because changes in bulk inventory (total atmospheric CO<sub>2</sub>) are confused with changes in isotopic <sup>14</sup>C/C ratio typically expressed as  $\Delta^{14}\text{C}$ . In detail, the concentration of CO<sub>2</sub> in the atmosphere, when perturbed by anthropogenic emissions, largely depends on the net oceanic CO<sub>2</sub> uptake rate, and therefore on the bottleneck of the carbonate chemistry as explained in section 3 above, while any changes in the isotopic <sup>14</sup>C/C ratio of atmospheric CO<sub>2</sub> also depends on the gross gas exchange rates (e.g. see Joos et al., 1996; Naegler and Levin, 2006). Furthermore, the record of atmospheric radiocarbon is perturbed/depleted by the emission of <sup>14</sup>C-free CO<sub>2</sub> from fossil fuels — the so-called <sup>14</sup>C Suess effect (Suess, 1955; Köhler, 2016) — and points to much longer atmospheric adjustment time on the order of 100 years for 1985–2005 (Levin et al., 2010).

#### 4. Harde's generalization including approximations based on paleo reconstructions

Finally, Harde (2017) generalizes the results that incorrectly model the modern carbon cycle to draw, again, erroneous conclusions about the paleoclimate record (section 3.3 and Fig. 3). Here, again, various shortcomings invalidate the conclusions. The main ones are the following:

- Glacial temperature:** The assumed surface temperature change for glacial times of  $-8$  K is wrong, as the source cited (Petit et al., 1999) approximates not global temperature change, but that over East Antarctica. According to some recent studies the global temperature change at the last glacial maximum with respect to pre-industrial times was  $-4.0 \pm 0.8$  K (Annan and Hargreaves, 2013). It is furthermore not clear to the reader how the data points in Fig. 3 were generated with one data point for a temperature of 8, 10, 12, 14°C, while the underlying paleo data from the Vostok ice core contain several thousand data points of the last 420,000 years, also including periods which have been warmer than the preindustrial climate.
- Explaining paleo CO<sub>2</sub>:** Harde (2017) proposes that the complete glacial/interglacial change in CO<sub>2</sub> can be explained by a reduction in surface temperature. However, it is nowadays well established, that glacial/interglacial changes in atmospheric CO<sub>2</sub> cannot be explained by one single process (e.g. Köhler et al., 2005; Brovkin et al., 2007; Kohfeld and Ridgwell, 2009). Significant change in atmospheric CO<sub>2</sub> on glacial/interglacial timescales is expected from a rise in sea surface temperature, rising sea-level, reduced marine export production, and responses from carbonate compensation, together with changing land carbon storage (Kohfeld and Ridgwell, 2009). Although models still disagree on the contribution of individual processes, the common consensus is, that the glacial/interglacial rise in temperature (more precisely sea surface temperature) might be responsible for a rise in atmospheric CO<sub>2</sub> mixing ratio of 20–30 ppm. The arguments in Harde (2017) are rather vague, but also seem to assume, that the temperature change might also trigger a change in terrestrial carbon storage. This concept would therefore need to have higher terrestrial carbon storage in cold periods, that might then be released during deglacial warming. However, the glacial terrestrial carbon storage is nowadays found to be smaller (not larger) in glacial times than during the pre-industrial period (Ciais et al., 2012). So, again evidence contradicts what is needed to support the concept set forth in Harde (2017). One might now argue, that all mentioned processes vary synchronously with temperature, and therefore the chosen temperature-dependency in Harde (2017) might be a possible simplification. However, this would largely ignore the complexity of the carbon cycle-climate system including the fact that the paleorecords also contain interglacial periods with higher than preindustrial global surface temperature, but similar atmospheric CO<sub>2</sub> concentrations, such as the last interglacial about 130,000 years ago (e.g. Bakker et al., 2013; Past Interglacials Working Group of PAGES, 2016).
- Paleo CO<sub>2</sub> data:** Furthermore, Harde (2017) argues that due to distortion and diffusion the CO<sub>2</sub> data from ice cores are rather imprecise leading to large error bars for CO<sub>2</sub> shown in Fig. 3 and to 20–30 ppm lower values than reconstructions based on fossil stomata. It must be clarified that ice core based CO<sub>2</sub> perfectly overlaps with the instrumental measurements of atmospheric CO<sub>2</sub> which started in 1958 and we therefore see no support for the contention that they might be biased to lower values (e.g. Ahn et al., 2012; Rubino et al., 2013; Köhler et al., 2017). Furthermore, short term variations seen in stomata-based CO<sub>2</sub> during the Holocene have been heavily criticized, and when averaged for known enclosure characteristics of gas bubbles in ice cores have not been confirmed in ice core-based records (e.g. Ahn et al., 2014; Köhler et al., 2015).



Together, the scientific arguments are in favor of the ice core-based CO<sub>2</sub> records and not of that based on fossil stomata.

- (d) **Impact of strong dependency of CO<sub>2</sub> on surface temperature:** Finally, the dependency of the atmospheric CO<sub>2</sub> concentration on the surface temperature, as depicted in Harde's Fig. 3 and his Eq. (17) would imply — as also discussed in Harde (2017) — a large contribution of CO<sub>2</sub> from the oceans for rising temperatures (but also some contributions from land). A large CO<sub>2</sub> outgassing contribution from the oceans, however, is at odds with the atmospheric oxygen records (e.g. see Keeling and Shertz, 1992).

## 5. Citations and some details on IPCC

Harde (2017) cites throughout various parts of the contribution of Working Group 1 (physical science basis) to the IPCC-AR5 (Stocker et al., 2013) and proposes *alternative views* on the impact of anthropogenic CO<sub>2</sub> emissions and related global temperature increase. This approach is inappropriate because it fails to address the actual underlying literature of the IPCC-AR5. The IPCC summarizes the state of the art in the peer-reviewed literature. Hence neither the residence time nor the adjustment time are assumptions or interpretations of the IPCC-AR5, but robust outcomes of the underlying science, which is summarized in the report. In presenting *alternative concepts* to the view presented in the IPCC-AR5 the article of Harde (2017) ignores and is in contradiction to the state of knowledge in the field, most obviously in the conflation of residence and adjustment times.

Some of the citations in Harde (2017) are inappropriate, including video presentations, and do not meet the standards of the peer-reviewed literature.

Further, Harde (2017) cites two papers (Essenhigh, 2009; Humlum et al., 2013) that were subject to highly critical peer-reviewed comments (Cawley, 2011; Kern and Leuenberger, 2013; Masters and Benestad, 2013; Richardson, 2013), none of which are referenced in Harde (2017). In fact, Harde (2017) repeats many of the same arguments that have already been refuted.

## 6. Conclusion

In conclusion, Harde (2017) does not provide an *alternative view* of the carbon cycle, but uses a too simplistic approach, that is based on invalid assumptions, and which leads to flawed results for anthropogenic carbon in the atmosphere. We suggest that the paper be withdrawn by the author, editor or publisher due to fundamental errors in the understanding of the carbon cycle.

## Acknowledgments

We thank Gavin Cawley for language editing and some stimulating discussions, Gavin Schmidt for starting this discussion on realclimate.org, and Fortunat Joos and two anonymous reviewers for their feedbacks.

## References

- Ahn, J., Brook, E.J., Buizert, C., 2014. Response of atmospheric CO<sub>2</sub> to the abrupt cooling event 8200 years ago. *Geophys. Res. Lett.* 41, 604–609. <http://dx.doi.org/10.1002/2013GL058177>.
- Ahn, J., Brook, E.J., Mitchell, L., Rosen, J., McConnell, J.R., Taylor, K., Etheridge, D., Rubino, M., 2012. Atmospheric CO<sub>2</sub> over the last 1000 years: a high-resolution record from the West Antarctic Ice Sheet (WAIS) divide ice core. *Glob. Biogeochem. Cycles* 26, GB2027. <http://dx.doi.org/10.1029/2011GB004247>.
- Annan, J.D., Hargreaves, J.C., 2013. A new global reconstruction of temperature changes at the Last Glacial Maximum. *Clim. Past* 9, 367–376. <http://dx.doi.org/10.5194/cp-9-367-2013>.
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven, G., Montenegro, A., Tokos, K., 2009. Atmospheric lifetime of fossil fuel carbon dioxide. *Annu. Rev. Earth Planet. Sci.* 37, 117–134. <http://dx.doi.org/10.1146/annurev.earth.031208.100206>.
- Archer, D., Khesghi, H., Maier-Reimer, E., 1997. Multiple timescales for neutralization of

- fossil fuel CO<sub>2</sub>. *Geophys. Res. Lett.* 24, 405–408. <http://dx.doi.org/10.1029/97GL00168>.
- Bakker, P., Stone, E.J., Charbit, S., Gröger, M., Krebs-Kanzow, U., Ritz, S.P., Varma, V., Khon, V., Lunt, D.J., Mikolajewicz, U., Prange, M., Renssen, H., Schneider, B., Schulz, M., 2013. Last interglacial temperature evolution – a model inter-comparison. *Clim. Past* 9, 605–619. <http://dx.doi.org/10.5194/cp-9-605-2013>.
- Brovkin, V., Ganopolski, A., Archer, D., Rahmstorf, S., 2007. Lowering of glacial atmospheric CO<sub>2</sub> in response to changes in oceanic circulation and marine biogeochemistry. *Paleoceanography* 22, PA4202. <http://dx.doi.org/10.1029/2006PA001380>.
- Cawley, G.C., 2011. On the Atmospheric residence time of anthropogenically sourced carbon dioxide. *Energy Fuel* 25, 5503–5513. <http://dx.doi.org/10.1021/ef200914u>.
- Ciais, P., Tagliabue, A., Cuntz, M., Bopp, L., Scholze, M., Hoffmann, G., Lourantou, A., Harrison, S.P., Prentice, I.C., Kelley, D.I., Koven, C., Piao, S.L., 2012. Large inert carbon pool in the terrestrial biosphere during the Last Glacial Maximum. *Nat. Geosci.* 5, 74–79. <http://dx.doi.org/10.1038/ngeo1324>.
- Dickson, A.G., Sabine, C.L., Christian, J.R., 2007. *Guide to Best Practices for Ocean CO<sub>2</sub> Measurements*. In: PICES Special Publication 3.
- Dlugokencky, E., Lang, P., Mund, J., Crotwell, A., Crotwell, M., Thoning, K., 2016. Atmospheric Carbon Dioxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1968–2015, Version: 2016-08-30. [ftp://aftp.cmdl.noaa.gov/data/trace\\_gases/co2/flask/surface/](ftp://aftp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/) Path:
- Eide, M., Olsen, A., Ninnemann, U.S., Eldevik, T., 2017. A global estimate of the full oceanic <sup>13</sup>C Suess effect since the preindustrial. *Glob. Biogeochem. Cycles* 31, 515–534. <http://dx.doi.org/10.1002/2016GB005472>.
- Essenhigh, R.H., 2009. Potential dependence of global warming on the residence time (RT) in the atmosphere of anthropogenically sourced carbon dioxide. *Energy Fuel* 23, 2773–2784. <http://dx.doi.org/10.1021/ef800581r>.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, F., Fung, I., Gnanadesikan, J., John, J., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H.D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A.J., Yoshikawa, C., Zeng, N., 2006. Climate-carbon cycle feedback analysis: results from the C<sup>4</sup>MIP model inter-comparison. *J. Clim.* 19, 3337–3353. <http://dx.doi.org/10.1175/JCLI3800.1>.
- Gruber, N., Gloor, M., Mikaloff Fletcher, S.E., Doney, S.C., Dutkiewicz, S., Follows, M.J., Gerber, M., Jacobson, A.R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S.A., Sarmiento, J.L., Takahashi, T., 2009. Oceanic sources, sinks, and transport of atmospheric CO<sub>2</sub>. *Glob. Biogeochem. Cycles* 23, GB1005. <http://dx.doi.org/10.1029/2008GB003349>.
- Gruber, N., Sarmiento, J.L., Stocker, T.F., 1996. An improved method for detecting anthropogenic CO<sub>2</sub> in the oceans. *Glob. Biogeochem. Cycles* 10, 809–837. <http://dx.doi.org/10.1029/96GB01608>.
- Harde, H., 2017. Scrutinizing the carbon cycle and CO<sub>2</sub> residence time in the atmosphere. *Glob. Planet. Chang.* 152, 19–26. <http://dx.doi.org/10.1016/j.gloplacha.2017.02.009>.
- Hauck, J., Hoppema, M., Bellerby, R.G.J., Völker, C., Wolf-Gladrow, D., 2010. Data-based estimation of anthropogenic carbon and acidification in the Weddell Sea on a decadal timescale. *J. Geophys. Res.* 115, C03004. <http://dx.doi.org/10.1029/2009JC005479>.
- Heimann, M., Maier-Reimer, E., 1996. On the relations between the oceanic uptake of CO<sub>2</sub> and its carbon isotopes. *Glob. Biogeochem. Cycles* 10, 89–110. <http://dx.doi.org/10.1029/95GB03191>.
- Houghton, J.T., Jenkins, G.J., Ephraums, J.J., 1990. *Climate Change - The IPCC Scientific Assessment*. Cambridge University Press, Cambridge, UK.
- Humlum, O., Stordahl, K., Solheim, J.-E., 2013. The phase relation between atmospheric carbon dioxide and global temperature. *Glob. Planet. Chang.* 100, 51–69. <http://dx.doi.org/10.1016/j.gloplacha.2012.08.008>.
- Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., Liddicoat, S., Lindsay, K., Reick, C.H., Roelandt, C., Segsneider, J., Tjiputra, J., 2013. Twenty-first-century compatible CO<sub>2</sub> emissions and airborne fraction simulated by CMIP5 earth system models under four representative concentration pathways. *J. Clim.* 26, 4398–4413. <http://dx.doi.org/10.1175/JCLI-D-12-00554.1>.
- Joos, F., Bruno, M., Fink, R., Siegenthaler, U., Stocker, T.F., Le Quéré, C., Sarmiento, J.L., 1996. An efficient and accurate representation of complex oceanic and biospheric models of anthropogenic carbon uptake. *Tellus* 48B, 397–417. <http://dx.doi.org/10.1034/j.1600-0889.1996.t01-2-00006.x>.
- Joos, F., Roth, R., Fuglestedt, J.S., Peters, G.P., Enting, I.G., von Bloh, W., Brovkin, V., Burke, E.J., Eby, M., Edwards, N.R., Friedrich, T., Frölicher, T.L., Halloran, P.R., Holden, P.B., Jones, C., Kleinen, T., Mackenzie, F.T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segsneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., Weaver, A.J., 2013. Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.* 13, 2793–2825. <http://dx.doi.org/10.5194/acp-13-2793-2013>.
- Keeling, R.F., Shertz, S.R., 1992. Seasonal and interannual variations in atmospheric oxygen and implications for the global carbon cycle. *Nature* 358, 723–727. <http://dx.doi.org/10.1038/358723a0>.
- Kern, Z., Leuenberger, M., 2013. Comment on "The phase relation between atmospheric carbon dioxide and global temperature" Humlum et al. [*Glob. Planet. Chang.* 100: 51–69]. Isotopes ignored. *Glob. Planet. Chang.* 109, 1–2. <http://dx.doi.org/10.1016/j.gloplacha.2013.07.002>.
- Kohfeld, K.E., Ridgwell, A., 2009. Glacial-interglacial variability in atmospheric CO<sub>2</sub>. In: Saltzman, E., Le Quéré, C. (Eds.), *Surface Ocean–Lower Atmospheres Processes*. volume 187 of *Geophysical Monograph Series*. AGU, Washington D.C., pp. 251–286. <http://dx.doi.org/10.1029/2008GM000845>.
- Köhler, P., 2016. Using the Suess effect on the stable carbon isotope to distinguish the future from the past in radiocarbon. *Environ. Res. Lett.* 11, 124016. <http://dx.doi.org/10.1088/1748-7598/11/12/124016>.

- <http://dx.doi.org/10.1088/1748-9326/11/12/124016>.
- Köhler, P., Fischer, H., Munhoven, G., Zeebe, R.E., 2005. Quantitative interpretation of atmospheric carbon records over the last glacial termination. *Glob. Biogeochem. Cycles* 19, GB4020. <http://dx.doi.org/10.1029/2004GB002345>.
- Köhler, P., Fischer, H., Schmitt, J., Brook, E.J., Marcott, S.A., 2015. Comment on "Synchronous records of  $p\text{CO}_2$  and  $\Delta^{14}\text{C}$  suggest rapid, ocean-derived  $p\text{CO}_2$  fluctuations at the onset of the Younger Dryas" by Steinthorsdottir et al. *Quat. Sci. Rev.* 107, 267–270. <http://dx.doi.org/10.1016/j.quascirev.2014.09.024>.
- Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T.F., Fischer, H., 2017. A 156 kyr smoothed history of the atmospheric greenhouse gases  $\text{CO}_2$ ,  $\text{CH}_4$ , and  $\text{N}_2\text{O}$  and their radiative forcing. *Earth Syst. Sci. Data* 9, 363–387. <http://dx.doi.org/10.5194/essd-9-363-2017>.
- Landschützer, P., Gruber, N., Bakker, D.C.E., 2016. Decadal variations and trends of the global ocean carbon sink. *Glob. Biogeochem. Cycles* 30, 1396–1417. <http://dx.doi.org/10.1002/2015GB005359>.
- Lauvset, S.K., Key, R.M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F.F., Suzuki, T., Watelet, S., 2016. A new global interior ocean mapped climatology: the  $1^\circ \times 1^\circ$  GLODAP version 2. *Earth Syst. Sci. Data* 8, 325–340. <http://dx.doi.org/10.5194/essd-8-325-2016>.
- Le Quéré, C., Andrew, R.M., Canadell, J.G., Sitch, S., Korsbakken, J.I., Peters, G.P., Manning, A.C., Boden, T.A., Tans, P.P., Houghton, R.A., Keeling, R.F., Alin, S., Andrews, O.D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L.P., Ciais, P., Currie, K., Delire, C., Doney, S.C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A.K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J.R., Metz, N., Millero, F., Monteiro, P.M.S., Munro, D.R., Nabel, J.E.M.S., Nakaoka, S.-I., O'Brien, K., Olsen, A., Omar, A.M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B.D., Sutton, A.J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I.T., van der Werf, G.R., Viovy, N., Walker, A.P., Wiltshire, A.J., Zaehle, S., 2016. Global Carbon Budget 2016. *Earth Syst. Sci. Data* 8, 605–649. <http://dx.doi.org/10.5194/essd-8-605-2016>.
- Levin, I., Naegler, T., Kromer, B., Diehl, M., Francey, R.J., Gomez-Pelaez, A.J., Steele, L.P., Wagenbach, D., Weller, R., Worthy, D.E., 2010. Observations and modelling of the global distribution and long-term trend of atmospheric  $^{14}\text{C}$ . *Tellus B* 62, 26–46. <http://dx.doi.org/10.1111/j.1600-0889.2009.00446.x>.
- Lord, N.S., Ridgwell, A., Thorne, M.C., Lunt, D.J., 2016. An impulse response function for the long tail of excess atmospheric  $\text{CO}_2$  in an Earth system model. *Glob. Biogeochem. Cycles* 30, 2–17. <http://dx.doi.org/10.1002/2014GB005074>.
- Masters, T., Benestad, R., 2013. Comment on "The phase relation between atmospheric carbon dioxide and global temperature". *Glob. Planet. Chang.* 106, 141–142. <http://dx.doi.org/10.1016/j.gloplacha.2013.03.010>.
- Naegler, T., Levin, I., 2006. Closing the global radiocarbon budget 1945–2005. *J. Geophys. Res.* 111, D12311. <http://dx.doi.org/10.1029/2005JD006758>.
- Oeschger, H., Siegenthaler, U., Schotterer, U., Gugelmann, A., 1975. A box diffusion model to study the carbon dioxide exchange in nature. *Tellus* 27, 168–192. <http://dx.doi.org/10.1111/j.2153-3490.1975.tb01671.x>.
- Orr, J.C., Epitalon, J.-M., Gattuso, J.-P., 2015. Comparison of ten packages that compute ocean carbonate chemistry. *Biogeosciences* 12, 1483–1510. <http://dx.doi.org/10.5194/bg-12-1483-2015>.
- Past Interglacials Working Group of PAGES, 2016. Interglacials of the last 800,000 years. *Rev. Geophys.* 54, 162–219. <http://dx.doi.org/10.1002/2015RG000482>.
- Petit, J.R., Jouzel, J., Raynaud, D., Barkov, N.I., Barnola, J.-M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V.M., Legrand, M., Lipenkov, V.Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E., Stievenard, M., 1999. Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature* 399, 429–436. <http://dx.doi.org/10.1038/20859>.
- Revelle, R., Suess, H.E., 1957. Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric  $\text{CO}_2$  during the past decades. *Tellus* 9, 18–27. <http://dx.doi.org/10.1111/j.2153-3490.1957.tb01849.x>.
- Richardson, M., 2013. Comment on "The phase relation between atmospheric carbon dioxide and global temperature" by Humlum, Stordahl and Solheim. *Glob. Planet. Chang.* 107, 226–228. <http://dx.doi.org/10.1016/j.gloplacha.2013.03.011>.
- Rubino, M., Etheridge, D.M., Trudinger, C.M., Allison, C.E., Battle, M.O., Langenfelds, R.L., Steele, L.P., Curran, M., Bender, M., White, J.W.C., Jenk, T.M., Blunier, T., Francey, R.J., 2013. A revised 1000-year atmospheric  $\delta^{13}\text{C}$ - $\text{CO}_2$  record from Law Dome and South Pole, Antarctica. *J. Geophys. Res.: Atmos.* 118, 8482–8499. <http://dx.doi.org/10.1002/jgrd.50668>.
- Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., Wanninkhof, R., Wong, C.S., Wallace, D.W.R., Tilbrook, B., Millero, F.J., Peng, T.-H., Kozyr, A., Ono, T., Rios, A.F., 2004. The oceanic sink for anthropogenic  $\text{CO}_2$ . *Science* 305, 367–371. <http://dx.doi.org/10.1126/science.1097403>.
- Stocker, T.F., Qin, D., Tignor, G.-K.P.M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P., 2013. IPCC, 2013: Climate Change 2013: The Physical Science Basis. In: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Suess, H.E., 1955. Radiocarbon concentration in modern wood. *Science* 122, 415–417. <http://dx.doi.org/10.1126/science.122.3166.415-a>.
- Volk, T., Hoffert, M.I., 1985. Ocean carbon pumps: analysis of relative strengths and efficiencies in ocean-driven atmospheric  $\text{CO}_2$  changes. In: Sundquist, E.T., Broecker, W.S. (Eds.), *The carbon cycle and atmospheric  $\text{CO}_2$ : natural variations Archean and present*. volume 32 of Geophysical Monograph. American Geophysical Union, Washington, D.C., USA, pp. 99–110. <http://dx.doi.org/10.1029/GM032p0099>.
- Zeebe, R.E., Wolf-Gladrow, D.A., 2001.  *$\text{CO}_2$  in Seawater: Equilibrium, Kinetics, Isotopes*. volume 65 of Elsevier Oceanography Book Series Elsevier Science Publishing, Amsterdam, The Netherlands.